

REFORESTATION WITH MAJOR BEE FOOD TREES IN EL SALVADOR

Marieke SANDKER

PROMABOS, Aportado Postal 1882, Centro de Gobierno, San Salvador, El Salvador
E-mail: mariekesandker@hotmail.com

Abstract

The preliminary results of a trial plantation with major bee food trees in El Salvador (Central America) will be shortly discussed. The 20 neotropical bee food trees will be presented and their characteristics for which they are selected in the reforestation will be highlighted. The trees are selected for the stingless bee *Melipona beecheii*, and planted on the land of beekeepers by the project "PROMABOS". The aim of the reforestation is increasing the honey production of *M. beecheii*, thus enhancing its survival and augmenting the income of local beekeepers. PROMABOS has included the reforestation program in the project activities because deforestation is the main hurdle in beekeeping with stingless bees in Central America. In El Salvador, only 2-5% of the land is covered with primary forest, around 30% with degraded woodland.

Introduction

In this article, the selection of major bee forage trees for reforestation near beecolonies is discussed. The reforestation is aiming the development of stingless beekeeping in El Salvador, Central America, as part of the stingless beekeeping project PROMABOS. The selection criteria for these plant species and conditions for the reforestation are reviewed, and the major bee forage trees and shrubs are listed with a brief review of their properties.

Stingless Beekeeping in El Salvador: PROMABOS

The native bee species in the tropical Americas are stingless bees (Apidae, Meliponinae). These bees vary strongly in size, honey production and honey quality. One of the most appreciated species for its honey is *Melipona beecheii*, a stingless bee endemic to Central America with about the same size as the honeybee (*Apis mellifera*). It was already kept by the Mayas in pre-hispanic times for its highly appreciated honey, which is produced in lower quantities than in honeybee colonies but distinguishes in its taste and more medicinal properties are subscribed to it, and therefore it sells at a better price. Owing to their particular swarming behaviour and the requirements for nest-sites, *M. beecheii* is very vulnerable to habitat fragmentation.

Massive deforestation has taken place in the tropical Americas and El Salvador is left only with 5 to 10% of its original vegetation (FAO, 2000). *M. beecheii* is in danger of extinction and in El Salvador it is kept only in certain regions in the North. Owing to its specific buzz-pollination in particular, this *Melipona* species seems important for the pollination and conservation of some of the indigenous trees and plants.

PROMABOS is a project aiming at the development of stingless beekeeping, specially the keeping of *M. beecheii*, in the region of La Palma, North-West of El Salvador. A major aspect of this project is a reforestation program with bee forage trees. These trees will be planted on the land of beekeepers with stingless bees in the region with the aim to increase the honey production of their *Melipona* colonies. For this reason, the project has inventoried the flora present in the region and is applying pollen studies to investigate what plant species are of greater importance in the diet of *M. beecheii*. On the plants listed in the inventory, information of different studies is gathered about their importance in the diet of *M. beecheii* and *A. mellifera* and major bee forage plants are selected for the reforestation program. In order to create a bee forage forest, 10 important nectariferous or polliferous trees forming the canopy are selected and for the understorey 10 important small trees or shrubs. These canopy and understorey trees are planted in trial plantations in May and June 2003. The mixing of species has the advantage of fast flowering of the fast growing shrubs and small trees while the larger trees develop and of creating a bee food understorey with the shade resistant shrubs; the disadvantage of mixed planting is that the faster growing species can take away light for the lower developing ones. The development of the mixed plantation under different ecological conditions will be monitored and accordingly a new selection will be made with the most promising species and recommendations on their planting and management will be provided.

The Region of La Palma

The region of La Palma is located in the North-West of El Salvador, close to the border with Honduras and Guatemala. The beekeepers with stingless bees are found approximately between 600 and

1300 meters above sealevel. The vegetation on the lower parts is a leafshedding tropical dry forest, and at the higher altitudes we find a tropical mountain cloud forest with pine (*Pinus occarpa*), oak (*Quercus spp*) and *Liquidambar styraciflua* as characteristic species.

Importance Plants in the Diet of Melipona beecheii

Different information sources have been reviewed to learn whether the plant species in the inventory are of importance for nectar and/or pollen supply to *M beecheii*. Since there exist only few studies on this stingless bee, also information on utilization of the plant species by *A mellifera* is included. Having about the same size, it could be that they have a lot of overlap in their diets.

The studies reviewed are based on different methods: from interviews with beekeepers to palynological studies of pollen taken from the worker bees returning from the field. In many studies the methods or source of the information is not provided, so one does not know how reliable the information really is. E.g., when a bee is seen to visit a flower, this does not necessarily imply it is collecting nectar or pollen from it; one literature source can be stated in four different studies making the reader mistakenly believe that the species is recommended by four independent studies. To prevent these misunderstandings and provide more transparent information, a scheme is made with detailed information of the method used to identify the plant species collected by *Melipona beecheii*, *Melipona sp* or *Apis mellifera* when this was known or retraceable by the author.

Scheme 1

Summary table of the different studies on major bee food plants

The different existing methods of palynological research on pollen mosters are given in the first four columns. The mosters are taken from the pollen or honey storage pots in the nests (stored pollen and stored honey), from the curbiculae of the worker bee returning from the field collecting pollen (pollen loads) and from the body or the nectar in the honey stomach of the worker bee returning from the field collecting nectar (nectar loads). The other method is field observation on the flower. The final column contains studies claiming the importance of the plant species for bees but without the method or results on which this information is based.

	Bee species	Stored pollen	Stored honey	Pollen loads	Nectar loads	Observation on flower			Unknown method		
						Nectar	Pollen	?	Nectar	Pollen	?
<i>Acacia angustissima</i>	Mb							33*			
	Am								10, 29, 30, 31	30	
<i>Anacardium occidentale</i>	Mb								3		
	Msp				1						
	Am			16					9, 10, 13, 19, 27, 30, 31	9, 19, 27, 30, 31	
<i>Andira inermis</i>	Mb							34	3		
	Am							34	3, 9, 10, 18, 19, 30		6
<i>Bixa orellana</i>	Mb									3	
	Msp	2		20a	1	5				19	
	Am						10			3, 19, 29	6
<i>Bursera simaruba</i>	Mb								3	3	
	Msp								19	19	
	Am		28	23, 24		10			3, 7, 19, 29, 31	3, 7, 19, 29, 30	
<i>Byrsonima crassifolia</i>	Msp										3
	Am					10			3, 9, 19, 29, 31	9, 29	
<i>Cassia grandis</i>	Mb										3
	Am								3, 10, 19	3	
<i>Cedrela odorata</i>	Mb			12						3	
	Am								3, 10, 29, 30, 31	3, 19	6
<i>Cochlospermum vitifolium</i>	Mb									3	
	Am								9, 10, 19, 26, 30, 31	3, 7, 10, 19, 26, 29, 30, 31	6, 26d, g, i, j
<i>Cordia alliodora</i>	Mb			12							
	Am		11	11, 24, 26				34	3, 7, 10, 13, 14, 19, 26, 27, 29, 30, 31	3, 19, 26, 27	6, 26a, b, c, f, g



<i>Croton reflexifolius</i>	Mb							33*, 34			
<i>Gliricidia sepium</i>	Am			26		10		34	9, 19, 26, 27, 29, 30, 31, 32	26, 31	26a, c, e, h, j
<i>Inga vera</i>	Am			24					9, 10, 19		
<i>Liquidambar styraciflua</i>	Mb							33*			
<i>Persea americana</i>	Mb							33*			
	Am					10			9, 13, 18, 19, 27, 29, 31	9, 19, 27	6
<i>Pithecelobium dulce</i>	Am								3, 9, 10, 19, 28, 29, 30, 31	3, 9, 19, 29, 30	
<i>Psidium guajava</i>	Mb							17	3	3	
	Msp	20b	20b	4, 22, 25						19	
	Am		11, 15	8, 11, 22				17	3, 9, 10, 13, 19, 27, 31	3, 9, 10, 19, 27	
<i>Spondias purpurea</i>	Am								3, 10, 29, 30, 31		19
<i>Tabebuia rosea</i>	Am								3, 10, 19, 26 (a, d, g) 29, 30, 31	3	
<i>Vernonia patens</i>	Mb							17, 33*			
	Am			24				17, 34	10, 19, 30	19	

* Since it is difficult to distinguish *Apis mellifera* and *Melipona beecheii* in the field, this information provided by the beekeepers with stingless bees is not very "species reliable".

References:	Locality:	References:	Locality:
1 Absy et al (1980)	Brazil (Manaus)	23 Roubik et al (1986)	Panama (Central)
2 Absy et al (1984)	Brazil (The Amazonas)	24 Sánchez-Chaves (1999)	Costa Rica (Guanacaste)
3 Arce et al (2001)	information from different countries	25 Sommeijer et al (1983)	Trinidad
4 Bootsma MC (1987?)	Trinidad	26 Stephen-Lobo (1999)	Costa Rica
5 Brantjes (1981)	Brazil (Pará)	26a Bently and Elias (1983) in 26 ?	
6 Chandrasekharan et al (1996)	source locality information unknown	26b Crane (1990) in 26	?
7 Chemas and Rico-Gray (1991)	Mexico (Yucatán)	26c Espina (1984) in 26	?
8 Cortopassi and Ramalho (1988)	Brazil (São Paulo)	26d Frankie (1983) in 26	?
9 Crane et al (1984)	information from different countries	26e Kerkvliet et al (1991) in 26	?
10 Espina and Ordetx (1983)	information from different countries	26f Martínez (1993) in 26	?
11 Girón-VanderHuck (1996)	Colombia	26g Palacios (1987) in 26	?
12 Landaverde-Parada (2003)	Costa Rica (Alajuela)	26h Ramalho (1990) in 26	?
13 Laurence (1973)	Trinidad	26i Roubik and Moreno (1991) in 26	?
14 León and Poveda (1999)	source locality information unknown	26j Vinson (1987) in 26	?
15 Lobreau-Callen et al (1986)	West-Africa (Togo and Benin)	27 Svensson (1991)	information from different countries
16 Magalhães-Freitas (1991)	Brazil	28 Villanueva (1994)	Mexico (Yucatán)
17 Marroquín-Juarez (1994)	Costa Rica	29 Villegas-Durán et al (1998)	Mexico (Yucatán)
18 Niembro-Rocas (1990)	source locality information unknown	30 Villegas-Durán et al (2002a)	Mexico (Guerrero)
19 Nieuwstadt van (1994)	information from different countries	31 Villegas-Durán et al (2002b)	Mexico (Chiapas)
20 Ramalho et al (1990)	Brazil (São Paulo)	32 Woyke (1983)	El Salvador
20a Absy and Kerr (1977) in 20	?	33 Important according to stingless beekeepers	El Salvador (La Palma region)
20b Kleinert and Imperatriz (1987) in 20	?	34 Personal observation	El Salvador (La Palma region)
21 Ramalho et al (1989)	Brazil		
22 Rooy de (1981)	Trinidad		

When interpreting this scheme, it is important to know that the same plant species will perform differently under different ecological and climatic conditions and therefore the same species can be a major forage tree in one region, while of no significant importance to the bees in another region (BRUYN, 1997). These facts underline the importance of local information from beekeepers and local information of the palynological research.

For this local importance, two plant species are included on which no information was found in the reviewed studies but which are mentioned as important by the beekeepers in the region, namely *Liquidambar styraciflua* and *Croton reflexifolius*. The reason they are not mentioned in the reviewed studies is probably that they do not occur in the region of study or very rarely.

Other Selection Criteria for Reforestation Plants

For the reforestation program to be successful, the situation in the field implies other selection criteria for the major bee forage plants as well besides their importance to bees.

When planting trees on the farmer's land, their development is only guaranteed when the farmers protect them from fire and cattle. They will do this only if the tree is sufficiently appreciated by them. So,

preferably the species has other uses besides being a major bee forage tree. The most appreciated characteristics gained by interviews with the beekeepers are trees producing timber (*A. occidentale*, *A. inermis*, *C. odorata*, *C. alliodora*, *I. vera*, *L. styraciflua* and *T. rosea*), producing fruit (*A. occidentale*, *Byrsonima crassifolia*, *C. grandis*, *I. vera*, *P. americana*, *P. guajava* and *S. purpurea*) or ornamentals (*A. angustissima*, *A. inermis*, *B. orellana*, *C. grandis*, *C. vitifolium*, *L. styraciflua*, *T. rosea* and *V. patens*).

The plants have to be applicable on different scales: if a farmer has no terrain available for reforestation, small trees or shrubs can be planted on the limited space near the bee hives where there are no possibilities for planting large growing trees. Furthermore, small trees and shrubs will give flowers shortly after planting, so in the bee forage forest they function as the "pioneers" in bee food provision. Large trees will be planted in the large-scale reforestation since their full-grown canopy gives a high quantity of flowers. The shrubs to small trees in the list are *A. angustissima*, *A. occidentale*, *B. orellana*, *B. crassifolia*, *C. reflexifolius*, *G. sepium*, *P. guajava*, *S. purpurea* and *V. patens*. The medium sized to large trees in the list are: *A. inermis*, *B. simaruba*, *C. grandis*, *C. odorata*, *C. vitifolium*, *C. alliodora*, *I. vera*, *L. styraciflua*, *P. americana*, *P. dulce* and *T. rosea*.

The tree has to develop well and have a high nectar flow under the local conditions. For this reason, trees are selected which already naturally occur in the region and in the trial plantations their development on different altitudes, different soils and different light conditions will be monitored. Furthermore, their nectar flow and pollen provision for *M. beecheii* will be investigated. Depending on their performance under the local conditions, a new selection will be made and for the following years the trees will be planted where they perform best.

Ultimately, the PROMABOS project aims at managing the natural forest and therefore only native species will be planted (species native to the tropical Americas) as not to disturb the natural composition of the forest.

The last criteria for the reforestation program to have an effect on the stingless bees is that the trees will be planted within the flight range of *M. beecheii* and preferably as close as possible. The degradation of importance with the distance to the beehive is not known so this should be investigated.

Summary of Criteria

Criterion 1: the trees must be primary source of pollen and / or nectar for *Melipona beecheii*.

Criterion 2: the trees must be appreciated by the landowners.

Criterion 3: in the selection we need large and small trees or shrubs to plant on different sized terrains.

Criterion 4: the tree must develop well and perform well as a bee forage source under the local conditions.

Criterion 5: the tree must be native to the tropical Americas.

Criterion 6: the trees must be planted near the beehives, at least within the flight range but preferable as close as possible.

REFERENCES

- Absy ML, Braga-Bezerra E and Kerr WE (1980), Plantas nectaríferas utilizadas por duas espécies de *Melipona* da Amazonia. *Acta Amazonica* 10 (2), pp271-281
- Absy ML, Camargo JMF, Kerr WE & Andrade de-Miranda IP (1984), Espécies de plantas visitadas por Meliponinae (Hymenoptera: Apoidea) para colecta de pólen na região do médio Amazonas. *Rev. Brasil. Biol.* 44 (2), pp 227-237
- Arce HG, Sánchez LA, Slaa J, Sánchez-Vindas PE, Ortiz AM, Veen JW Van and Sommeijer MJ (2001), Árboles melíferos nativos de Mesoamérica. PRAM, Heredia, pp 207
- Bootsma MC (1989), Vergelijkend onderzoek naar stuifmeelbronnen van *Melipona trinitatis* en *Apis mellifera* in Trinidad, West-Indies. Student thesis UU, Utrecht, pp 21
- Brantjes NBM (1981), Nectar and the pollination of bread fruit, *Artocarpus altilis* (Moraceae). *Acta Botanica Neerlandica* 30 (5/6), pp 345-352
- Bruyn Cde (1997), Practical beekeeping. The Crowood Press, Ramsbury, pp 288
- Chandrasekharan C, Frisk T & Campos-Roasio J (1996), Desarrollo de productos forestales no madereros en América Latina y el Caribe, website: <http://www.fao.org/docrep/t2360s/t2360s0h.htm>, reviewed 04/2003
- Chemas A and Rico-Gray V (1991), Apiculture and management of associated vegetation by the maya of Tixcacaltuyub, Yucatán, Mexico. *Agroforestry Systems* 13, pp 13-25
- Cortopassi-Laurino M and Ramalho M (1988), Pollen harvest by Africanized *Apis mellifera* and *Trigona spinipes* in São Paulo botanical and ecological views. *Apidologie* 19 (1), pp 1-24
- Crane E., Walker P & Day R (1984), Directory of important world honey resources. IBRA, London, pp 384
- Espina Perez D and Ordetx Ros GS (1983), Flora Apícola Tropical Costa Rica. Editorial Tecnológica de Costa Rica, Costa Rica, pp 406
- FAO (2000) FAO-Forestry. Website: http://www.fao.org/forestry/fo/country/index.jsp?lang_id=1&geo_id=173, reviewed 5-2003
- Girón-VanderHuck M (1996), Recolección de polen y néctar por *Apis mellifera* en algunas especies de plantas silvestres y cultivadas del municipio de Salgar (Antioquia). In: Melitopalínología. Litografía Luz, Calarcá Colombia, pp 83
- Landaverde-Parada VL (2003), Informe de capacitación en palinología. Work report PRAM, Costa Rica, pp 46
- Laurence GA (1973), Some bee plants of Trinidad. *Journal of the Agricultural Society of Trinidad and Tobago* 73, pp100-101

- León J and Poveda LJ (1999), Nombres comunes de las plantas en Costa Rica. Editorial Fundación UNA, Costa Rica, pp 870
- Lobreau-Callen D, Darchen R and Le Thomas A (1986), Apport de la palynologie a la connaissance des relations abeilles/plantes en savanes arborées du Togo et du Bénin. *Apidologie* 17 (4), pp 279-306
- Magalhães-Freitas B (1991), Potencial da caatinga para produção de polen e néctar para a exploração apícola. Dissertation Ceara, Brazil, pp 1-114
- Marroquín-Juarez AE (1994), Traslape de recursos florales entre abejas sin aguijón (Meliponinae) y *Apis mellifera* (L.). Work report PRAM, Heredia, pp 20
- Niembro Rocas A (1990), Árboles y arbustos útiles de México. Editorial Limusa, México DF, pp 206
- Nieuwstadt M van (1994), CARABIS Central American Apibotanical Information System. Computer database PRAM, Heredia-Costa Rica, pp 453
- Ramalho A, Kleinert-Giovannini A, and Imperatriz-Fonseca VL (1990), Important bee plants for stingless bees (*Melipona* and *Trigonini*) and Africanized honeybees (*Apis mellifera*) in neotropical habitats: a review. *Apidologie* 21, pp 469-488
- Ramalho M, Kleinert-Giovannini A and Imperatriz-Fonseca (1989), Utilization of floral resources by species of *Melipona* (Apidae, Meliponinae): floral preferences. *Apidologie* 20, pp 185-195
- Rooy B de (1981), Aspecten van het fourageergedrag van angelloze bijen op Trinidad met nadruk op de herkomst van het verzamelde stuifmeel. Student thesis UU, Utrecht, pp 44
- Roubik DW, Moreno EJ, Vergara C and Wittman D (1986), Sporadic food competition with the African honeybee: projected impact on neotropical social bees. *Journal of Tropical Ecology* 2, pp 97-111
- Sánchez-Chaves LA (1999), Floral preferences of the native bee *Tetragonisca angustula* (Apidae: meliponinae) and the Africanized honeybee *Apis mellifera* (Apidae: Apinae) spectrum, diversity and overlap of the pollen diet. Student thesis PRAM-UU, Heredia – Utrecht, pp 44
- Sommeijer MJ, Rooy GA de, Punt W and Bruijn LLM de (1983), A comparative study of foraging behavior and pollen resources of various stingless bees (Hym., Meliponinae) and honeybees (Hym., apinae) in Trinidad, West-Indies. *Apidologie* 14(3), pp 205-224
- Stephen-Lobo EP (1999), Factores que condicionan la importancia de las plantas apícolas. In: Student thesis PRAM, Heredia, pp 47
- Svensson B (1991), Bees and trees. Working paper 183 Swedish University of Agricultural Sciences, Uppsala, pp 80
- Villanueva-G R (1994), Nectar sources of European and Africanized honey bees (*Apis mellifera* L.) in the Yucatán Peninsula, Mexico. *Journal of Apicultural Research* 33 (1), pp 44-58
- Villegas-Durán G, Cajero-Avelar S, Bolaños-Medina A, Miranda-Sánchez JA, Pérez-Lara MA et al (1998), Flora nectarífera y polinífera de la Península de Yucatán. SAGARPA, Mexico DF, pp 126
- Villegas-Durán G, Bolaños-Medina A, Miranda-Sánchez JA & González Quintero U (2002a), Flora nectarífera y polinífera en el estado de Guerrero. SAGARPA, Mexico DF, pp 126
- Villegas-Durán G, Bolaños-Medina A, Miranda-Sánchez JA & Zenón-Abarca AJ (2002b), Flora nectarífera y polinífera en el estado de Chiapas. SAGARPA, Mexico DF, pp 164
- Woyke HJ (1983), La apicultura en El Salvador. FAO-MAG, San Salvador, pp16