

Royal jelly and propolis prevent development of insulin resistance in type 2 diabetic animal models.

Hiromu Kawasaki 1), Yoshito Zamami 1), Toshihiro Koyama 1) , Hiroki Fujiwara 1), Shima Takaki-Doi 2), Tomoki Tatefuji 2) and Ken Hashimoto 2)

1) Department of Clinical Pharmaceutical Science, Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama University, and 2) Institute for Bee Products & Health Science, Yamada Apiculture Center, Inc.

Honeybee products such as royal jelly (RJ) and propolis (PPL) are known to contain excellent nutrition and have a variety of biological activities. The present study was designed to investigate the effects of RJ and PPL on insulin resistance (hyperinsulinemia) in fructose-drinking rats (FDR; insulin resistance and type 2 diabetic animal model) and Otsuka Long-Evans Tokushima Fatty (OLETF; type 2 diabetic animal model) rats. Male Wistar rats (6 week-old) received 15% fructose solution in drinking water for 8 weeks. FDR showed significant increases in plasma levels of insulin and triglyceride, Homeostasis Model Assessment ratio (HOMA-R, an index of insulin resistance), and systolic blood pressure, but not blood glucose levels, when compared with control rats. RJ (100 and 300 mg/kg/day, p.o.) or PPL (Brazilian propolis extract; 100 and 300 mg/kg/day, p.o.) treatment for 8 weeks significantly decreased the plasma levels of insulin and triglyceride, HOMA-R, without affecting blood glucose or total cholesterol levels and tended to lower systolic blood pressure. In another insulin resistance model OLETF rats, RJ (30 and 300 mg/kg/day, p.o.) or PPL (100 and 300 mg/kg/day, p.o.) treatment for 4 weeks tended to decrease systolic blood pressure and significantly decreased serum level of insulin and HOMA-R. These results suggest that RJ and PPL could be an effective functional food to prevent insulin resistance associated with the development of hypertension.